

WI 15 ROCK POWDER

PROPERTIES

Rock powder, with fortifying, rebalancing and strengthening action for plants and soils. It is suitable for any type of cultivation and soil.

APPLICATION

WIDDAR WI 15 ROCK POWDER is recommended to **assist propagation via CUTTINGS**. Use at a rate of 10 g(0.35 oz.) of powder diluted in 200 ml(6.75 fluid oz.) of clean water. The cuttings should be left in the solution as follows:

- **Herbaceous cuttings:** 15 – 60 minutes
- **Softwood cuttings:** 75 – 120 minutes
- **Hardwood cuttings:** 100 – 180 minutes

STORAGE: Keep the product dry and well sealed. Keep out of the reach of children.

AVAILABLE IN: 3 kg (6.6 lbs), 5 kg (11 lbs), 10 kg (22 lbs), 25 kg (55 lbs) packs.

For information on the correct application in specific soil and climate conditions and cultivation methods, please contact our technical support office.

USAGE

Rock powder can be diluted in two ways:

- Pour the powder in the sprayer barrel in this dosage: 1 kg/ha (2.2 lb./2.5 acres) to which you add 200/250 litres (50-65 gal.) of water. After about 30 minutes, the product will be ready to be sprayed.
- In the event of clogged sprayer nozzles, or excessive wear of the pump's bearings, pour the powder into a bucket, add 10/15 (2.64-4 gallons) litres of water, stir 2-3 times and after about 30 minutes, pour the solution into the sprayer barrel using a suitable filter. Lastly, add the rest of the water.

Please note that the diluted solution does not expire when kept in a tightly closed tank.

NOTICE

Please note that the usage of the specific products on the plants and the soil needs to always be used within the whole **Trinium** agricultural method, otherwise they will not be effective. The products must also be contained and applied using barrels and pumps free of any residue of non-Widdar products. Barrels must be thoroughly washed with caustic soda or, alternatively, treated internally and externally with **WI 21 ROCK POWDER** fortifier.

If it rains during the 3 hours following the application of Widdar products the effectiveness of the treatment is reduced by about 20%. If it rains within 2 hours the products will be ineffective and will have to be reapplied.

AVERAGE MINERALOGICAL ANALYSIS		OTHER CHARACTERISTICS	
QUARTZ	20-40%	MOISTURE	<5%
CALCITE	5-30%	pH	6,5-7%
SODIUM FELDSPAR	5-35%		
POTASSIUM FELDSPAR	5-30%		
OTHER	5-20%		

* Method licensed from EUREKA Coop.

Certified by **bioagri^{inputs}cert**
IFOAM
ACCREDITED

This product is included at point 5) and listed in Annex I of Italian Ministerial Decree 18354 of 27/11/2009 setting out the arrangements for the implementation of EC Regulations no. 834/2007, no. 889/2008, no. 1235/2008 and later additions and amendments concerning organic production. Product not subject to authorization by the Italian Health Ministry as per Ministerial Decree 18/07/2018, n. 6793 and Presidential Decree 28/02/2012 no. 55.